

**RECEIVED
CENTRAL FAX CENTER****FEB 20 2007**
Attorney's Docket: 2000DE463
Serial No.: 10/025,065
Group: 1716REMARKS

The Office Action mailed October 20, 2006, has been carefully considered together with each of the references cited therein. The amendments and remarks presented herein are believed to be fully responsive to the Office Action. The amendments made herein are fully supported by the application as originally filed. No new matter has been added. Accordingly, reconsideration of the present Application in view of the above amendments and following remarks is respectfully requested.

Claim Status

Claims 1 and 3-8 are pending in this application. By this Amendment, Applicants have amended claims 5 through 8.

Claim Objections

Claim 6 stands rejected to as the Office finds it depends from a cancelled claim. Claim 2 has been amended, changing its dependency from claim 2 to claim 1, thereby overcoming the instant objection.

Claim Rejections Under 35 USC § 103

Claims 1 and 3-8 stand rejected under 35 USC § 103(a) as being unpatentable over U.S. Patent No. 6,489,395 to Loffler et al. in view of the article entitled "Hydrafresh with the Right Polymer" by Dr. Rainer Weihofen, Clariant, February 2001. This rejection is respectfully traversed.

Applicants hereby perfect the priority claim of December 23, 2000 by attaching herewith a copy of an English translation of the certified copy of the foreign priority document along with a statement that the translation of the certified copy is accurate. The certified copy of DE 10065046.5 was mailed to the Office on December 19, 2001.

BEST AVAILABLE COPY

Attorney's Docket: 2000DE463
Serial No.: 10/025,065
Group: 1716

Basis for the claimed subject matter can be found in the attached English translation, *inter alia*, as follows:

Claim 1: page 2, lines 19-26; page 3, lines 23-25; the examples beginning on page 28 and ending on page 31 and page 32, claim 1;

Claim 3: page 3, lines 20 through page 4 line 5;

Claim 4: page 32, claim 4;

Claim 5: page 32, claim 5;

Claim 6: page 3, lines 23-25;

Claim 7: page 32, claim 5; and

Claim 8: page 32, claim 5.

In view of the above, Applicants courteously believe that priority to DE 10065046.5 of December 23, 2000 has been perfected. In consequence, the article entitled "Hydrafresh with the Right Polymer" by Dr. Rainer Weihofen, Clariant, published February 2001, does not constitute prior art under § 102 against the instant Application. Applicants, therefore, respectfully request reconsideration and withdrawal of the § 103 rejection.

Attorney's Docket: 2000DE463
Serial No.: 10/025,065
Group: 1716

In view of the forgoing amendments and remarks, the present Application is believed to be in condition for allowance, and reconsideration of it is requested. If the Examiner disagrees, she is requested to contact the attorney for Applicants at the telephone number provided below.

Respectfully submitted,




Anthony A. Bisulca
Attorney for Applicant
Registration No. 40,913

(CUSTOMER NUMBER 25,255)

Clariant Corporation
Industrial Property Department
4000 Monroe Road
Charlotte, North Carolina 28205
Phone: (704) 331-7151
Fax: (704) 331-7707

RWS Group Ltd, of Europa House, Marsham Way, Gerrards Cross, Buckinghamshire, England, hereby declares that, to the best of its knowledge and belief, the following document, prepared by one of its translators competent in the art and conversant with the English and German languages, is a true and correct translation of the accompanying German Patent Application No. 100 65 046.5 filed on 23 December 2000.

Signed this 1st day of February 2007



C. E. SITCH

Acting Managing Director

For and on behalf of RWS Group Ltd

RECEIVED
CENTRAL FAX CENTER

FEB 20 2007

FEDERAL REPUBLIC OF GERMANY

[Eagle crest]

Priority Certificate

DE 100 65 046.5

for the filing of a Patent Application

File Reference: 100 65 046.5

Filing date: 23 December 2000

Applicant/Proprietor: Clariant International Ltd., Muttentz/CH

First Applicant: Beiersdorf AG, 20253 Hamburg/DE

Title: O/W emulsions containing one or more ammonium
acryloyldimethyltaurate/vinylpyrrolidone copolymers

IPC: A 61 K 8/02, A 61 Q 99/00, A 61 Q 1/10,
A 61 Q 17/00, A 61 Q 19/00, A 61 K 8/81,
A 61 K 8/92, A 61 K 8/30, A 61 K 47/00

The attached documents are a correct and accurate reproduction of the parts of the documents submitted on 23 December 2000 for this Application notwithstanding any colour changes due to the copying process.

Munich, 17 January 2007

German Patent and Trademark Office

The President

PP

[Seal of the German Patent
and Trademark Office]

[signature]

Remus

Beiersdorf Aktiengesellschaft
Hamburg

5

Description

O/W emulsions containing one or more ammonium
acryloyldimethyltaurate/vinylpyrrolidone copolymers

10

The present invention relates to cosmetic and dermatological preparations of the oil-in-water type, to processes for their preparation, and to their use for cosmetic and medicinal purposes.

15

The human skin is man's largest organ and performs a number of vital functions. Having an average area of about 2 m² in adults, it has a prominent role as a protective and sensory organ. The purpose of this organ is to transmit and avert mechanical, thermal, actinic, chemical and biological stimuli. In addition, it has an important role as a regulatory and target organ in human metabolism.

20

The main aim of skin care in the cosmetics sense is to strengthen or restore the skin's natural function as a barrier against environmental influences (e.g. dirt, chemicals, microorganisms) and against the loss of endogenous substances (e.g. water, natural fats, electrolytes), and also to assist its horny layer in its natural regeneration ability in cases of existing damage.

25

If the barrier properties of the skin are impaired, increased resorption of toxic or allergenic substances or attack by microorganisms may result, leading to toxic or allergic skin reactions.

30

Another aim of skin care is to compensate for the loss by the skin of sebum and water caused by daily washing. This is particularly important if the natural regeneration ability is inadequate. Furthermore, skin care products should protect against environmental influences, in particular against sun and wind, and delay skin aging.

PAWI/2000/200-195.doc

Medicinal topical compositions usually comprise one or more medicaments in an effective concentration. For the sake of simplicity, in order to clearly distinguish between cosmetic and medicinal use and corresponding products, reference is made to the legal provisions in the Federal Republic of Germany (e.g. Cosmetics Directive, Foods and Drugs Act).

5

Emulsions are generally understood as meaning heterogeneous systems which consist of two liquids, which are usually referred to as phases, which are immiscible or miscible with one another only to a limited extent. In an emulsion, one of the two liquids is dispersed in the form of very fine droplets in the other liquid.

10

If the two liquids are water and oil and oil droplets are very finely dispersed in water, this is an oil-in-water emulsion (O/W emulsion, e.g. milk). The basic character of an O/W emulsion is determined by the water. In the case of a water-in-oil emulsion (W/O emulsion, e.g. butter), the principle is reversed, the basic character being determined here

15

by the oil.

The aim was to remedy these shortcomings.

20

Surprisingly, these objects are achieved by cosmetic or dermatological emulsions of the oil-in-water type, comprising


- (i) up to 90% by weight of a water phase,
- (ii) up to 20% by weight of a lipid phase, based on the total weight of the preparations,
- (iii) up to 5% by weight of one or more emulsifiers,
- (iv) also comprising up to 5% by weight of one or more ammonium acryloyldimethyl-

25

taurates/vinylpyrrolidone copolymers.

30

According to the invention, the ammonium acryloyldimethyltaurates/vinylpyrrolidone copolymer(s) have the empirical formula $[C_7H_{16}N_2SO_4]_n [C_6H_9NO]_m$, corresponding to a statistical structure as follows



Preferred species for the purposes of the present invention are filed in Chemical Abstracts under the registry numbers 58374-69-9, 13162-05-5 and 88-12-0 and are obtainable under the trade name Aristoflex® AVC from Clariant GmbH.

5

It had therefore not been foreseen by the person skilled in the art that the preparations according to the invention

- would have better effectiveness as moisture-donating preparations,
- would be easier to formulate,
- 10 - would better promote skin smoothing,
- would be characterized by better care action,
- would better serve as vehicles for cosmetic and medicinal-dermatological active ingredients
- would have better sensory properties, such as, for example, the ability to be
- 15 - distributed on the skin or the ability to be absorbed into the skin,
- would have higher stability against decomposition in oil and water phases and
- would be characterized by better biocompatibility

than the preparations of the prior art.

- 20 The preparations according to the invention thus represent an enrichment of the prior art.

The lipid content of the preparations obtainable according to the invention can advantageously be varied from 0.5% by weight to 20% by weight, preferably from 5 to

- 25 10% by weight, where the results achieved are equally favorable. In the case of

freedom from lipid, no emulsion is present, but rather a system which should most appropriately be referred to as an emulsifier gel.

Preparations according to the invention preferably comprise up to 7.5% by weight of a lipid phase. Preparations according to the invention particularly advantageously
5 comprise up to 6% by weight of a lipid phase. Preparations according to the invention particularly preferably comprise 2 to 4% by weight of a lipid phase, in particular approximately 3% by weight, in each case based on the total weight of the preparations.

10

The lipid phase of the cosmetic or dermatological emulsions according to the invention can advantageously be chosen from the following group of substances:

- mineral oils, mineral waxes
- oils, such as triglycerides of capric or of caprylic acid, and also natural oils such
15 as, for example, castor oil;
- fats, waxes and other natural and synthetic fatty substances, preferably esters of fatty acids with alcohols of low carbon number, e.g. with isopropanol, propylene glycol or glycerol, or esters of fatty alcohols with alkanolic acids of low carbon number or with fatty acids;
- 20 - alkyl benzoates;
- silicone oils, such as dimethylpolysiloxanes, diethylpolysiloxanes, diphenylpolysiloxanes and mixed forms thereof.

The oil phase of the emulsions of the present invention is advantageously chosen from
25 the group of esters of saturated and/or unsaturated, branched and/or unbranched alkanecarboxylic acids having a chain length of from 3 to 30 carbon atoms and saturated and/or unsaturated, branched and/or unbranched alcohols having a chain length of from 3 to 30 carbon atoms, from the group of esters of aromatic carboxylic acids and saturated and/or unsaturated, branched and/or unbranched alcohols having
30 a chain length of from 3 to 30 carbon atoms. Such ester oils can then advantageously be chosen from the group consisting of isopropyl myristate, isopropyl palmitate, isopropyl stearate, isopropyl oleate, n-butyl stearate, n-hexyl laurate, n-decyl oleate, isooctyl stearate, isononyl stearate, isononyl isononanoate, 2-ethylhexyl palmitate, 2-ethylhexyl laurate, 2-hexyldecyl stearate, 2-octyldecyl palmitate, oleyl oleate, oleyl

erucate, erucyl oleate, erucyl erucate, and synthetic, semisynthetic and natural mixtures of such esters, e.g. jojoba oil.

In addition, the oil phase can advantageously be chosen from the group of branched
5 and unbranched hydrocarbons and hydrocarbon waxes, of silicone oils, of dialkyl
ethers, the group of saturated or unsaturated, branched or unbranched alcohols, and
the fatty acid triglycerides, namely the triglycerol esters of saturated and/or
unsaturated, branched and/or unbranched alkanecarboxylic acids having a chain
length of from 8 to 24, in particular 12 - 18 carbon atoms. The fatty acid triglycerides
10 can, for example, advantageously be chosen from the group of synthetic,
semisynthetic and natural oils, e.g. olive oil, sunflower oil, soybean oil, groundnut oil,
rapeseed oil, almond oil, palm oil, coconut oil, palm kernel oil and the like.

Any mixtures of such oil and wax components can also be used advantageously for
15 the purposes of the present invention. It may also in some instances be advantageous
to use waxes, for example cetyl palmitate, as the sole lipid component of the oil phase.


The oil phase is advantageously chosen from the group consisting of 2-ethylhexyl
isostearate, octyldodecanol, isotridecyl isononanoate, isoeicosane, 2-ethylhexyl
20 cocoate, C₁₂₋₁₅-alkyl benzoate, caprylic/capric triglyceride, dicaprylyl ether.

Particularly advantageous mixtures are those of C₁₂₋₁₅-alkyl benzoate and 2-ethylhexyl
isostearate, mixtures of C₁₂₋₁₅-alkyl benzoate and isotridecyl isononanoate, and
mixtures of C₁₂₋₁₅-alkyl benzoate, 2-ethylhexyl isostearate and isotridecyl
25 isononanoate.


Of the hydrocarbons, paraffin oil, squalane and squalene are to be used
advantageously for the purposes of the present invention.

30 The oil phase can advantageously also have a content of cyclic or linear silicone oils,
or consist entirely of such oils, although it is preferable to use an additional content of
other oil phase components apart from the silicone oil or the silicone oils. Such
silicones or silicone oils may be in the form of monomers, which are generally
characterized by structural elements, as follows:

6




Linear silicones having two or more siloxyl units which are to be used advantageously according to the invention are generally characterized by structural elements, as follows:



where the silicon atoms can be substituted by identical or different alkyl radicals and/or aryl radicals, which are shown here in general terms by the radicals $R_1 - R_4$ (that is to say the number of different radicals is not necessarily limited to 4). m can assume values from 2 - 200 000.

Cyclic silicones to be used advantageously according to the invention are generally characterized by structural elements, as follows



where the silicon atoms can be substituted by identical or different alkyl radicals and/or aryl radicals, which are shown here in general terms by the radicals $R_1 - R_4$ (that is to say the number of different radicals is not necessarily limited to 4). n can assume values from 3/2 to 20. Fractions for n take into consideration that uneven numbers of siloxyl groups may be present in the cycle.

Advantageously, cyclomethicone (e.g. decamethylcyclopentasiloxane) is used as the silicone oil to be used according to the invention. However, other silicone oils are also to be used advantageously for the purpose of the present invention, for example

undecamethylcyclotrisiloxane, polydimethylsiloxane, poly(methylphenylsiloxane), cetyl-dimethicone, behenoxydimethicone.

Also advantageous are mixtures of cyclomethicone and isotridecyl isononanoate, and
5 those of cyclomethicone and 2-ethylhexyl isostearate.

It is, however, also advantageous to choose silicone oils of similar constitution to the
above-described compounds whose organic side chains are derivatized, for example
polyethoxylated and/or polypropoxylated. These include, for example, polysiloxane-
10 polyalkyl-polyether copolymers, such as cetyl-dimethicone copolyol, (cetyl-dimethicone
copolyol (and) polyglyceryl-4-isostearate (and) hexyl laurate).

Also particularly advantageous are mixtures of cyclomethicone and isotridecyl
isononanoate, and of cyclomethicone and 2-ethylhexyl isostearate.

15

The aqueous phase of the preparations according to the invention optionally
advantageously comprises alcohols, diols or polyols of low carbon number, and ethers
thereof, preferably ethanol, isopropanol, propylene glycol, glycerol, ethylene glycol,
ethylene glycol monoethyl or monobutyl ether, propylene glycol monomethyl,
20 monoethyl or monobutyl ether, diethylene glycol monomethyl or monoethyl ether and
analogous products, and also alcohols of low carbon number, e.g. ethanol,
isopropanol, 1,2-propanediol, glycerol, and, in particular, one or more thickeners which
can advantageously be chosen from the group consisting of silicon dioxide and
aluminum silicates.

25

Preparations according to the invention in the form of emulsions advantageously
comprise, in particular, one or more hydrocolloids. These hydrocolloids can
advantageously be chosen from the group of gums, polysaccharides, cellulose
derivatives, phyllosilicates, polyacrylates and/or other polymers.

30

The gums include saps from plants or trees which harden in the air and form resins, or
extracts from aquatic plants. From this group, for the purposes of the present
invention, gum arabic, carob flour, tragacanth, karaya, guar gum, pectin, gellan gum,

carrageen, agar, algins, chondrus, xanthan gum, for example, can be chosen advantageously.

Also advantageous is the use of derivatized gums, such as, for example,
5 hydroxypropyl guar (Jaguar® HP 8).

The polysaccharides and polysaccharide derivatives include, for example, hyaluronic acid, chitin and chitosan, chondroitin sulfates, starch and starch derivatives.

10 The cellulose derivatives include, for example, methylcellulose, carboxymethylcellulose, hydroxyethylcellulose, hydroxypropylmethylcellulose.

The phyllosilicates include naturally occurring and synthetic clay earths, such as, for example, montmorillonite, bentonite, hectorite, laponite, magnesium aluminum
15 silicates such as Veegum®. These can be used as such or in modified form, such as, for example, stearylalkonium hectorite.

In addition, silica gels can also be used advantageously.

20 The polyacrylates include, for example, Carbopol grades from Goodrich (Carbopol 980, 981, 1382, 5984, 2984, EDT 2001 or Pemulen TR2).

The polymers include, for example, polyacrylamides (Seppigel 305), polyvinyl alcohols, PVP, PVP/VA copolymers, polyglycols.

25

Preparation according to the invention in the form of emulsions comprise one or more emulsifiers. These emulsifiers can advantageously be chosen from the group of nonionic, anionic, cationic or amphoteric emulsifiers.

30 The nonionic emulsifiers include

- a) partial fatty acid esters and fatty acid esters of polyhydric alcohols and ethoxylated derivatives thereof (e.g. glyceryl monostearates, sorbitan stearates, glyceryl stearyl citrates, sucrose stearates)
- b) ethoxylated fatty alcohols and fatty acids

- c) ethoxylated fatty amines, fatty acid amides, fatty acid alkanolamides
- d) alkylphenol polyglycol ethers (e.g. Triton X).

The anionic emulsifiers include

- 5 a) soaps (e.g. sodium stearate)
- b) fatty alcohol sulfates
- c) mono-, di- and trialkyl phosphoric esters and ethoxylates thereof.

The cationic emulsifiers include

- 10 a) quaternary ammonium compounds with a long-chain aliphatic radical, e.g. distearyldimonium chloride.

The amphoteric emulsifiers include

- a) alkylaminoalkancarboxylic acids
- 15 b) betaines, sulfobetaines
- c) imidazoline derivatives.

In addition, there are naturally occurring emulsifiers, which include beeswax, wool wax, lecithin and sterols.

20

O/W emulsifiers can be advantageously chosen, for example, from the group of polyethoxylated or polypropoxylated or polyethoxylated and polypropoxylated products, e.g.:

- fatty alcohol ethoxylates,
- 25 - ethoxylated wool wax alcohols,
- polyethylene glycol ethers of the general formula $R-O-(CH_2-CH_2-O)_n-R'$,
- fatty acid ethoxylates of the general formula $R-COO-(CH_2-CH_2-O)_n-H$,
- etherified fatty acid ethoxylates of the general formula $R-COO-(CH_2-CH_2-O)_n-R'$,
- 30 - esterified fatty acid ethoxylates of the general formula $R-COO-(CH_2-CH_2-O)_n-C(O)-R'$,
- polyethylene glycol glycerol fatty acid esters,
- ethoxylated sorbitan esters,

10

- cholesterol ethoxylates,
- ethoxylated triglycerides,
- alkyl ether carboxylic acids of the general formula
 $R-O-(-CH_2-CH_2-O-)_n-CH_2-COOH$ and n is a number from 5 to 30,
- 5 - polyoxyethylene sorbitol fatty acid esters,
- alkyl ether sulfates of the general formula $R-O-(-CH_2-CH_2-O-)_n-SO_3-H$,
- fatty alcohol propoxylates of the general formula
 $R-O-(-CH_2-CH(CH_3)-O-)_n-H$,
- polypropylene glycol ethers of the general formula
 $R-O-(-CH_2-CH(CH_3)-O-)_n-R'$,
- 10 - propoxylated wool wax alcohols,
- etherified fatty acid propoxylates
 $R-COO-(-CH_2-CH(CH_3)-O-)_n-R'$,
- esterified fatty acid propoxylates of the general formula
 $R-COO-(-CH_2-CH(CH_3)-O-)_n-C(O)-R'$,
- 15 - fatty acid propoxylates of the general formula
 $R-COO-(-CH_2-CH(CH_3)-O-)_n-H$,
- polypropylene glycol glycerol fatty acid esters,
- propoxylated sorbitan esters,
- 20 - cholesterol propoxylates,
- propoxylated triglycerides,
- alkyl ether carboxylic acids of the general formula
 $R-O-(-CH_2-CH(CH_3)-O-)_n-CH_2-COOH$,
- alkyl ether sulfates or the parent acids of these sulfates of the general formula
 $R-O-(-CH_2-CH(CH_3)-O-)_n-SO_3-H$,
- 25 - fatty alcohol ethoxylates/propoxylates of the general formula
 $R-O-X_n-Y_m-H$,
- polypropylene glycol ethers of the general formula
 $R-O-X_n-Y_m-R'$,
- 30 - etherified fatty acid propoxylates of the general formula
 $R-COO-X_n-Y_m-R'$,
- fatty acid ethoxylates/propoxylates of the general formula
 $R-COO-X_n-Y_m-H$.

According to the invention, particularly advantageous polyethoxylated or polypropoxylated or polyethoxylated and polypropoxylated O/W emulsifiers used are those chosen from the group of substances having HLB values of 11 - 18, very particularly preferably having HLB values of 14.5 - 15.5, provided the O/W emulsifiers have saturated radicals R and R'. If the O/W emulsifiers have unsaturated radicals R and/or R', or isoalkyl derivatives are present, then the preferred HLB value of such emulsifiers can also be lower or higher.

It is advantageous to choose the fatty alcohol ethoxylates from the group of ethoxylated stearyl alcohols, cetyl alcohols, cetylstearyl alcohols (cetearyl alcohols). Particular preference is given to:

polyethylene glycol(13) stearyl ether (steareth-13), polyethylene glycol(14) stearyl ether (steareth-14), polyethylene glycol(15) stearyl ether (steareth-15), polyethylene glycol(16) stearyl ether (steareth-16), polyethylene glycol(17) stearyl ether (steareth-17), polyethylene glycol(18) stearyl ether (steareth-18), polyethylene glycol(19) stearyl ether (steareth-19), polyethylene glycol(20) stearyl ether (steareth-20),

polyethylene glycol(12) isostearyl ether (isosteareth-12), polyethylene glycol(13) isostearyl ether (isosteareth-13), polyethylene glycol(14) isostearyl ether (isosteareth-14), polyethylene glycol(15) isostearyl ether (isosteareth-15), polyethylene glycol(16) isostearyl ether (isosteareth-16), polyethylene glycol(17) isostearyl ether (isosteareth-17), polyethylene glycol(18) isostearyl ether (isosteareth-18), polyethylene glycol(19) isostearyl ether (isosteareth-19), polyethylene glycol(20) isostearyl ether (isosteareth-20),

polyethylene glycol(13) cetyl ether (ceteth-13), polyethylene glycol(14) cetyl ether (ceteth-14), polyethylene glycol(15) cetyl ether (ceteth-15), polyethylene glycol(16) cetyl ether (ceteth-16), polyethylene glycol(17) cetyl ether (ceteth-17), polyethylene glycol(18) cetyl ether (ceteth-18), polyethylene glycol(19) cetyl ether (ceteth-19), polyethylene glycol(20) cetyl ether (ceteth-20),

polyethylene glycol(13) isocetyl ether (isoceteth-13), polyethylene glycol(14) isocetyl ether (isoceteth-14), polyethylene glycol(15) isocetyl ether (isoceteth-15), polyethylene

glycol(16) isocetyl ether (isoceteth-16), polyethylene glycol(17) isocetyl ether (isoceteth-17), polyethylene glycol(18) isocetyl ether (isoceteth-18), polyethylene glycol(19) isocetyl ether (isoceteth-19), polyethylene glycol(20) isocetyl ether (isoceteth-20),

5

polyethylene glycol(12) oleyl ether (oleth-12), polyethylene glycol(13) oleyl ether (oleth-13), polyethylene glycol(14) oleyl ether (oleth-14), polyethylene glycol(15) oleyl ether (oleth-15),

10 polyethylene glycol(12) lauryl ether (laureth-12), polyethylene glycol(12) isolauryl ether (isolaureth-12),

polyethylene glycol(13) cetylstearyl ether (cetareth-13), polyethylene glycol(14) cetylstearyl ether (cetareth-14), polyethylene glycol(15) cetylstearyl ether (cetareth-15), polyethylene glycol(16) cetylstearyl ether (cetareth-16), polyethylene glycol(17) cetylstearyl ether (cetareth-17), polyethylene glycol(18) cetylstearyl ether (cetareth-18), polyethylene glycol(19) cetylstearyl ether (cetareth-19), polyethylene glycol(20) cetylstearyl ether (cetareth-20).

20 It is also advantageous to choose the fatty acid ethoxylates from the following group:

polyethylene glycol(20) stearate, polyethylene glycol(21) stearate, polyethylene glycol(22) stearate, polyethylene glycol(23) stearate, polyethylene glycol(24) stearate, polyethylene glycol(25) stearate,

25

polyethylene glycol(12) isostearate, polyethylene glycol(13) isostearate, polyethylene glycol(14) isostearate, polyethylene glycol(15) isostearate, polyethylene glycol(16) isostearate, polyethylene glycol(17) isostearate, polyethylene glycol(18) isostearate, polyethylene glycol(19) isostearate, polyethylene glycol(20) isostearate, polyethylene glycol(21) isostearate, polyethylene glycol(22) isostearate, polyethylene glycol(23) isostearate, polyethylene glycol(24) isostearate, polyethylene glycol(25) isostearate,

30

polyethylene glycol(12) oleate, polyethylene glycol(13) oleate, polyethylene glycol(14) oleate, polyethylene glycol(15) oleate, polyethylene glycol(16) oleate, polyethylene

13

glycol(17) oleate, polyethylene glycol(18) oleate, polyethylene glycol(19) oleate, polyethylene glycol(20) oleate.

5 The ethoxylated alkyl ether carboxylic acid or salt thereof which can be used is advantageously sodium laureth-11 carboxylate.

Sodium laureth-14 sulfate can be used advantageously as alkyl ether sulfate.

10 An advantageous ethoxylated cholesterol derivative which can be used is polyethylene glycol(30) cholesteryl ether. Polyethylene glycol(25) soyasterol has also proven successful.

Ethoxylated triglycerides which can be advantageously used are polyethylene glycol(60) Evening Primrose glycerides.

15

It is also advantageous to choose the polyethylene glycol glycerol fatty acid esters from the group polyethylene glycol(20) glyceryl laurate, polyethylene glycol(21) glyceryl laurate, polyethylene glycol(22) glyceryl laurate, polyethylene glycol(23) glyceryl laurate, polyethylene glycol(6) glyceryl caprate, polyethylene glycol(20) glyceryl oleate, 20 polyethylene glycol(20) glyceryl isostearate, polyethylene glycol(18) glyceryl oleate/cocotate.

It is likewise favorable to choose the sorbitan esters from the group polyethylene glycol(20) sorbitan monolaurate, polyethylene glycol(20) sorbitan monostearate, 25 polyethylene glycol(20) sorbitan monoisostearate, polyethylene glycol(20) sorbitan monopalmitate, polyethylene glycol(20) sorbitan monooleate.

Advantageous W/O emulsifiers which can be used are: fatty alcohols having 8 to 30 carbon atoms, monoglycerol esters of saturated and/or unsaturated, branched and/or 30 unbranched alkanecarboxylic acids having a chain length of from 8 to 24, in particular 12 - 18, carbon atoms, diglycerol esters of saturated and/or unsaturated, branched and/or unbranched alkanecarboxylic acids having a chain length of from 8 to 24, in particular 12 - 18, carbon atoms, monoglycerol ethers of saturated and/or unsaturated, branched and/or unbranched alcohols having a chain length of from 8 to 24, in

14

- particular 12 - 18, carbon atoms, diglycerol ethers of saturated and/or unsaturated, branched and/or unbranched alcohols having a chain length of from 8 to 24, in particular 12 - 18, carbon atoms, propylene glycol esters of saturated and/or unsaturated, branched and/or unbranched alkanecarboxylic acids having a chain
- 5 length of from 8 to 24, in particular 12 - 18, carbon atoms, and sorbitan esters of saturated and/or unsaturated, branched and/or unbranched alkanecarboxylic acids having a chain length of from 8 to 24, in particular 12 - 18, carbon atoms.

- Particularly advantageous W/O emulsifiers are glyceryl monostearate, glyceryl
- 10 monoisostearate, glyceryl monomyristate, glyceryl monooleate, diglyceryl monostearate, diglyceryl monoisostearate, propylene glycol monostearate, propylene glycol monoisostearate, propylene glycol monocaprylate, propylene glycol monolaurate, sorbitan monoisostearate, sorbitan monolaurate, sorbitan monocaprylate, sorbitan monoisoleate, sucrose distearate, cetyl alcohol, stearyl
- 15 alcohol, arachidyl alcohol, behenyl alcohol, isobehenyl alcohol, selachyl alcohol, chimyl alcohol, polyethylene glycol(2) stearyl ether (steareth-2), glyceryl monolaurate, glyceryl monocaprate, glyceryl monocaprylate.

- The emulsions according to the invention can comprise dyes and/or color pigments.
- 20 The dyes and color pigments can be chosen from the corresponding positive list of the Cosmetics Directive or the EC list of cosmetic colorants. In most cases they are identical to the dyes approved for foods. Advantageous color pigments are, for example, titanium dioxide, mica, iron oxides (e.g. Fe_2O_3 , Fe_3O_4 , $\text{FeO}(\text{OH})$) and/or tin oxide. Advantageous dyes are, for example, carmine, Berlin blue, chrome oxide green,
- 25 ultramarine blue and/or manganese violet. It is particularly advantageous to choose the dyes and/or color pigments from the following list. The Colour Index Numbers (CIN) are taken from the *Rowe Colour Index, 3rd Edition, Society of Dyers and Colourists, Bradford, England, 1971*.

Chemical or other name	CIN	Color
Pigment Green	10006	green
Acid Green 1	10020	green
2,4-Dinitrohydroxynaphthalene-7-sulfonic acid	10316	yellow
Pigment Yellow 1	11680	yellow

15

Chemical or other name	CIN	Color
Pigment Yellow 3	11710	yellow
Pigment Orange 1	11725	orange
2,4-Dihydroxyazobenzene	11920	orange
Solvent Red 3	12010	red
1-(2'-Chloro-4'-nitro-1'-phenylazo)-2-hydroxynaphthalene	12085	red
Pigment Red 3	12120	red
Ceres red; Sudan red; Fat Red G	12150	red
Pigment Red 112	12370	red
Pigment Red 7	12420	red
Pigment Brown 1	12480	brown
4-(2'-Methoxy-5'-sulfodiethylamido-1'-phenylazo)-3-hydroxy-5"-chloro-2",4"-dimethoxy-2-naphthanilide	12490	red
Disperse Yellow 16	12700	yellow
1-(4-Sulfo-1-phenylazo)-4-aminobenzene-5-sulfonic acid	13015	yellow
2,4-Dihydroxyazobenzene-4'-sulfonic acid	14270	orange
2-(2,4-Dimethylphenylazo-5-sulfo)-1-hydroxynaphthalene-4-sulfonic acid	14700	red
2-(4-Sulfo-1-naphthylazo)-1-naphthol-4-sulfonic acid	14720	red
2-(6-Sulfo-2,4-xylylazo)-1-naphthol-5-sulfonic acid	14815	red
1-(4'-Sulfophenylazo)-2-hydroxynaphthalene	15510	orange
1-(2-Sulfo-4-chloro-5-carboxy-1-phenylazo)-2-hydroxynaphthalene	15525	red
1-(3-Methylphenylazo-4-sulfo)-2-hydroxynaphthalene	15580	red
1-(4',8')-Sulfonaphthylazo)-2-hydroxynaphthalene	15620	red
2-Hydroxy-1,2'-azonaphthalene-1'-sulfonic acid	15630	red
3-Hydroxy-4-phenylazo-2-naphthylcarboxylic acid	15800	red
1-(2-Sulfo-4-methyl-1-phenylazo)-2-naphthylcarboxylic acid	15850	red
1-(2-Sulfo-4-methyl-5-chloro-1-phenylazo)-2-hydroxynaphthalene-3-carboxylic acid	15865	red
1-(2-Sulfo-1-naphthylazo)-2-hydroxynaphthalene-3-carboxylic acid	15880	red
1-(3-Sulfo-1-phenylazo)-2-naphthol-6-sulfonic acid	15980	orange
1-(4-Sulfo-1-phenylazo)-2-naphthol-6-sulfonic acid	15985	yellow
Allura Red	16035	red
1-(4-Sulfo-1-naphthylazo)-2-naphthol-3,6-disulfonic acid	16185	red
Acid Orange 10	16230	orange
1-(4-Sulfo-1-naphthylazo)-2-naphthol-6,8-disulfonic acid	16255	red

Chemical or other name	CIN	Color
1-(4-Sulfo-1-naphthylazo)-2-naphthol-3,6,8-trisulfonic acid	16290	red
8-Amino-2-phenylazo-1-naphthol-3,6-disulfonic acid	17200	red
Acid Red 1	18050	red
Acid Red 155	18130	red
Acid Yellow 121	18690	yellow
Acid Red 180	18736	red
Acid Yellow 11	18820	yellow
Acid Yellow 17	18965	yellow
4-(4-Sulfo-1-phenylazo)-1-(4-sulfophenyl)-5-hydroxy-pyrazolone-3-carboxylic acid	19140	yellow
Pigment Yellow 16	20040	yellow
2,6-(4'-Sulfo-2'', 4''-dimethyl)bisphenylazo)-1,3-dihydroxybenzene	20170	orange
Acid Black 1	20470	black
Pigment Yellow 13	21100	yellow
Pigment Yellow 83	21108	yellow
Solvent Yellow	21230	yellow
Acid Red 163	24790	red
Acid Red 73	27290	red
2-[4'-(4''-Sulfo-1''-phenylazo)-7'-sulfo-1'-naphthylazo]-1-hydroxy-7-aminonaphthalene-3,6-disulfonic acid	27755	black
4'-[(4''-Sulfo-1''-phenylazo)-7'-sulfo-1'-naphthylazo]-1-hydroxy-8-acetylaminonaphthalene-3,5-disulfonic acid	28440	black
Direct Orange 34, 39, 44, 46, 60	40215	orange
Food Yellow	40800	orange
trans- β -Apo-8'-carotinaldehyde (C ₃₀)	40820	orange
trans-Apo-8'-carotinic acid (C ₃₀)-ethyl ester	40825	orange
Canthaxanthin	40850	orange
Acid Blue 1	42045	blue
2,4-Disulfo-5-hydroxy-4'-4''-bis(diethylamino)triphenylcarbinol	42051	blue
4-[(4-N-Ethyl-p-sulfobenzylamino)phenyl(4-hydroxy-2-sulfophenyl)(methylene)-1-(N-ethyl-N-p-sulfobenzyl)-2,5-cyclohexadienimine]	42053	green
Acid Blue 7	42080	blue
(N-Ethyl-p-sulfobenzylamino)phenyl(2-sulfophenyl)methylene-(N-ethyl-N-p-sulfobenzyl) $\Delta^{2,5}$ -cyclohexadienimine	42090	blue

Chemical or other name	CIN	Color
Acid Green 9	42100	green
Diethyldisulfobenzyl-di-4-amino-2-chloro-di-2-methyl-fuchsonimmonium	42170	green
Basic Violet 14	42510	violet
Basic Violet 2	42520	violet
2'-Methyl-4'-(N-ethyl-N-m-sulfobenzyl)amino-4''-(N-diethyl)amino-2-methyl-N-ethyl-N-m-sulfobenzylfuchsonimmonium	42735	blue
4'-(N-Dimethyl)amino-4''-(N-phenyl)aminonaphtho-N-dimethyl-fuchsonimmonium	44045	blue
2-Hydroxy-3,6-disulfo-4,4'-bisdimethylaminonaphtho-fuchsonimmonium	44090	green
Acid Red 52	45100	red
3-(2'-Methylphenylamino)-6-(2'-methyl-4'-sulfophenylamino)-9-(2''-carboxyphenyl)xanthenium salt	45190	violet
Acid Red 50	45220	red
Phenyl-2-oxyfluorone-2-carboxylic acid	45350	yellow
4,5-Dibromofluorescein	45370	orange
2,4,5,7-Tetrabromofluorescein	45380	red
Solvent Dye	45396	orange
Acid Red 98	45405	red
3',4',5',6'-Tetrachloro-2,4,5,7-tetrabromofluorescein	45410	red
4,5-Diodofluorescein	45425	red
2,4,5,7-Tetraiodofluorescein	45430	red
Quinophthalone	47000	yellow
Quinophthalonedisulfonic acid	47005	yellow
Acid Violet 50	50325	violet
Acid Black 2	50420	black
Pigment Violet 23	51319	violet
1,2-Dioxyanthraquinone, calcium-aluminum complex	58000	red
3-Oxypyrene-5,8,10-sulfonic acid	59040	green
1-Hydroxy-4-N-phenylaminoanthraquinone	60724	violet
1-Hydroxy-4-(4'-methylphenylamino)anthraquinone	60725	violet
Acid Violet 23	60730	violet
1,4-Di(4'-methylphenylamino)anthraquinone	61565	green
1,4-Bis(o-sulfo-p-toluidino)anthraquinone	61570	green

Chemical or other name	CIN	Color
Acid Blue 80	61585	blue
Acid Blue 62	62045	blue
N,N'-Dihydro-1,2,1',2'-anthraquinone azine	69800	blue
Vat Blue 6; Pigment Blue 64	69825	blue
Vat Orange 7	71105	orange
Indigo	73000	blue
Indigo-disulfonic acid	73015	blue
4,4'-Dimethyl-6,6'-dichlorothioindigo	73360	red
5,5'-Dichloro-7,7'-dimethylthioindigo	73385	violet
Quinacridone Violet 19	73900	violet
Pigment Red 122	73915	red
Pigment Blue 16	74100	blue
Phthalocyanine	74160	blue
Direct Blue 86	74180	blue
Chlorinated phthalocyanine	74260	green
Natural Yellow 6,19; Natural Red 1	75100	yellow
Bixin, Norbixin	75120	orange
Lycopene	75125	yellow
trans-alpha-, beta- and gamma-carotene	75130	orange
Keto- and/or hydroxyl derivatives of carotene	75135	yellow
Guanine or pearly agent	75170	white
1,7-Bis(4-hydroxy-3-methoxyphenyl)-1,6-heptadiene-3,5-dione	75300	yellow
Complex salt (Na, Al, Ca) of carminic acid	75470	red
Chlorophyll a and b; copper compounds of chlorophylls and chlorophyllins	75810	green
Aluminum	77000	white
Hydrated alumina	77002	white
Hydrous aluminum silicates	77004	white
Ultramarine	77007	blue
Pigment Red 101 and 102	77015	red
Barium sulfate	77120	white
Bismuth oxychloride and its mixtures with mica	77163	white
Calcium carbonate	77220	white
Calcium sulfate	77231	white
Carbon	77266	black

Chemical or other name	CIN	Color
Pigment black 9	77267	black
Carbo medicinalis vegetabilis	77268:1	black
Chromium oxide	77288	green
Chromium oxide, hydrous	77289	green
Pigment Blue 28, Pigment Green 14	77346	green
Pigment Metal 2	77400	brown
Gold	77480	brown
Iron oxides and hydroxides	77489	orange
Iron oxide	77491	red
Hydrated Iron oxide	77492	yellow
Iron oxide	77499	black
Mixtures of iron(II)- and iron(III)hexacyanoferrate	77510	blue
Pigment White 18	77713	white
Manganese ammonium diphosphate	77742	violet
Manganese phosphate; $Mn_3(PO_4)_2 \cdot 7 H_2O$	77745	red
Silver	77820	white
Titanium dioxide and its mixtures with mica	77891	white
Zinc oxide	77947	white
6,7-Dimethyl-9-(1'-D-ribityl)isoalloxazine, lactoflavine		yellow
Sugar coloring		brown
Capsanthin, capsorubin		orange
Betanin		red
Benzopyrylium salts, anthocyan		red
Aluminium, zinc, magnesium and calcium stearate		white
Bromothymol blue		blue
Bromocresol green		green
Acid Red 195		red

- It may also be favorable to choose as dye one or more substances from the following group: 2,4-dihydroxyazobenzene, 1-(2'-chloro-4'-nitro-1'-phenylazo)-2-hydroxynaphthalene, Ceres Red, 2-(4-sulfo-1-naphthylazo)-1-naphthol-4-sulfonic acid, calcium salt of 2-hydroxy-1,2'-azonaphthalene-1'-sulfonic acid, calcium and barium salts of 1-(2-sulfo-4-methyl-1-phenylazo)-2-naphthylcarboxylic acid, calcium salt of 1-(2-sulfo-1-naphthylazo)-2-hydroxynaphthalene-3-carboxylic acid, aluminum salt of 1-(4-sulfo-1-phenylazo)-2-naphthol-6-sulfonic acid, aluminum salt of 1-(4-sulfo-1-naphthylazo)-2-

naphthol-3,6-disulfonic acid, 1-(4-sulfo-1-naphthylazo)-2-naphthol-6,8-disulfonic acid, aluminum salt of 4-(4-sulfo-1-phenylazo)-1-(4-sulfophenyl)-5-hydroxypyrazolone-3-carboxylic acid, aluminum and zirconium salts of 4,5-dibromofluorescein, aluminum and zirconium salts of 2,4,5,7-tetrabromofluorescein, 3',4',5',6'-tetrachloro-
5 2,4,5,7-tetrabromofluorescein and its aluminum salt, aluminum salt of 2,4,5,7-tetraiodofluorescein, aluminum salt of quinophthalone disulfonic acid, aluminum salt of indigo disulfonic acid; red and black iron oxide (CIN: 77 491 (red) and 77 499 (black)), iron oxide hydrate (CIN: 77 492), manganese ammonium diphosphate and titanium dioxide.

10

Also advantageous are oil-soluble natural dyes, such as, for example, paprika extracts, β -carotene or cochénille.

Also advantageous for the purposes of the present invention are emulsions with a
15 content of pearlescent pigments. Preference is given in particular to the types of pearlescent pigments listed below:

1. Natural pearlescent pigments, such as, for example
 - "pearl essence" (guanine/hypoxanthin mixed crystals from fish scales) and
 - "mother of pearl" (ground mussel shells)
- 20 2. Monocrystalline pearlescent pigments, such as, for example, bismuth oxychloride (BiOCl)
3. Layer-substrate pigments: e.g. mica/metal oxide.

Bases for pearlescent pigments are, for example, pulverulent pigments or castor oil
25 dispersions of bismuth oxychloride and/or titanium dioxide, and bismuth oxychloride and/or titanium dioxide on mica. The luster pigment listed under CIN 77163, for example, is particularly advantageous.

Also advantageous are, for example, the following types of pearlescent pigment based on
30 mica/metal oxide:

Group	Coating/layer thickness	Color
Silver-white pearlescent pigments	TiO ₂ : 40 – 60 nm	Silver
Interference pigments	TiO ₂ : 60 – 80 nm	Yellow
	TiO ₂ : 80 – 100 nm	Red
	TiO ₂ : 100 – 140 nm	Blue
	TiO ₂ : 120 – 160 nm	Green
Color luster pigments	Fe ₂ O ₃	Bronze
	Fe ₂ O ₃	Copper
	Fe ₂ O ₃	Red
	Fe ₂ O ₃	Red-violet
	Fe ₂ O ₃	Red-green
	Fe ₂ O ₃	Black
Combination pigments	TiO ₂ / Fe ₂ O ₃	Gold shades
	TiO ₂ / Cr ₂ O ₃	Green
	TiO ₂ / Berlin blue	Deep blue
	TiO ₂ / carmine	Red

Particular preference is given, for example, to the pearlescent pigments obtainable from Merck under the trade names Timiron, Colorona or Dichrona.

- 5 The list of given pearlescent pigments is not of course intended to be limiting. Pearlescent pigments which are advantageous for the purposes of the present invention are obtainable by numerous methods known per se. For example, other substrates apart from mica can be coated with further metal oxides, such as, for example, silica and the like. SiO₂ particles coated with, for example, TiO₂ and Fe₂O₃
- 10 ("ronaspheres"), which are marketed by Merck and are particularly suitable for the optical reduction of fine lines, are advantageous.

- It can moreover be advantageous to dispense completely with a substrate such as mica. Particular preference is given to iron pearlescent pigments prepared without the
- 15 use of mica. Such pigments are obtainable, for example, under the trade name Sicopearl Kupfer 1000 from BASF.

- In addition, also particularly advantageous are effect pigments which are obtainable under the trade name Metasome Standard/Glitter in various colors (yellow, red, green,
- 20 blue) from Flora Tech. The glitter particles are present here in mixtures with various

auxiliaries and dyes (such as, for example, the dyes with the Colour Index (CI) Numbers 19140, 77007, 77289, 77491).

5 The dyes and pigments may be present either individually or in a mixture, and can be mutually coated with one another, different coating thicknesses generally giving rise to different color effects. The total amount of dyes and color-imparting pigments is advantageously chosen from the range from e.g. 0.1% by weight to 30% by weight, preferably from 0.5 to 15% by weight, in particular from 1.0 to 10% by weight, in each case based on the total weight of the preparations.

10

The gel creams according to the invention can, in particular, advantageously be used as eyeshadows.

15 Particularly advantageous preparations are also obtained when antioxidants are used as additives or active ingredients: According to the invention, the preparations advantageously comprise one or more antioxidants. Antioxidants which are favorable but which are nevertheless optional may be all antioxidants which are customary or suitable for cosmetic and/or dermatological applications.

20 It is also advantageous to add antioxidants to the preparations according to the invention. The antioxidants are advantageously chosen from the group consisting of amino acids (e.g. glycine, histidine, tyrosine, tryptophan) and derivatives thereof, imidazoles (e.g. urocanic acid) and derivatives thereof, peptides, such as D,L-carnosine, D-carnosine, L-carnosine and derivatives thereof (e.g. anserine), carotenoids, carotenes
25 (e.g. α -carotene, β -carotene, lycopene) and derivatives thereof, chlorogenic acid and derivatives thereof, lipoic acid and derivatives thereof (e.g. dihydrolipoic acid), aurothio-glucose, propylthiouracil and other thiols (e.g. thioredoxin, glutathione, cysteine, cystine, cystamine and the glycosyl, N-acetyl, methyl, ethyl, propyl, amyl, butyl and lauryl, palmitoyl, oleyl, γ -linoleyl, cholesteryl and glyceryl esters thereof) and salts thereof,
30 dilauryl thiodipropionate, distearyl thiodipropionate, thiodipropionic acid and derivatives thereof (esters, ethers, peptides, lipids, nucleotides, nucleosides and salts) and sulfoximine compounds (e.g. buthionine sulfoximines, homocysteine sulfoximine, buthionine sulfones, penta-, hexa-, heptathionine sulfoximine) in very low tolerated doses (e.g. pmol to μ mol/kg), and also (metal) chelating agents (e.g. α -hydroxy fatty
35 acids, palmitic acid, phytic acid, lactoferrin), α -hydroxy acids (e.g. citric acid, lactic acid,

malic acid), humic acid, bile acid, bile extracts, bilirubin, biliverdin, EDTA, EGTA and derivatives thereof, unsaturated fatty acids and derivatives thereof (e.g. γ -linolenic acid, linoleic acid, oleic acid), folic acid and derivatives thereof, ubiquinone and ubiquinol and derivatives thereof, vitamin C and derivatives (e.g. ascorbyl palmitate, Mg ascorbyl phosphate, ascorbyl acetate), tocopherols and derivatives (e.g. vitamin E acetate), vitamin A and derivatives (vitamin A palmitate) and conferyl benzoate of benzoin resin, rutinic acid and derivatives thereof, α -glycosylrutin, ferulic acid, furfurylidene-glucitol, carnosine, butylhydroxytoluene, butylhydroxyanisole, nordihydroguaiacic acid, nordihydroguaiaretic acid, trihydroxybutyrophenone, uric acid and derivatives thereof, mannose and derivatives thereof, zinc and derivatives thereof (e.g. ZnO, ZnSO₄), selenium and derivatives thereof (e.g. selenomethionine), stilbenes and derivatives thereof (e.g. stilbene oxide, trans-stilbene oxide), and the derivatives (salts, esters, ethers, sugars, nucleotides, nucleosides, peptides and lipids) of said active ingredients which are suitable according to the invention.

15

For the purposes of the present invention, oil-soluble antioxidants can be used particularly advantageously.

A surprising property of the present invention is that preparations according to the invention are very good vehicles for cosmetic or dermatological active ingredients into the skin, preferred active ingredients being antioxidants which are able to protect the skin against oxidative stress. Preferred antioxidants are vitamin E and derivatives thereof and vitamin A and derivatives thereof.

20

The amount of antioxidants (one or more compounds) in the preparations is preferably from 0.001 to 30% by weight, particularly preferably 0.05-20% by weight, in particular 1-10% by weight, based on the total weight of the preparation.

25

If vitamin E and/or derivatives thereof are the antioxidant or antioxidants, the respective concentrations thereof are advantageously chosen from the range 0.001 - 10% by weight, based on the total weight of the formulation.

30

If vitamin A or vitamin A derivatives or carotenes or derivatives thereof are used as the antioxidant or antioxidants, the respective concentrations thereof are advantageously chosen from the range 0.001 - 10% by weight, based on the total weight of the formulation.

35

The person skilled in the art is of course aware that cosmetic preparations are in most cases inconceivable without the customary auxiliaries and additives. The cosmetic and dermatological preparations according to the invention can, accordingly, also comprise
5 cosmetic auxiliaries, as are customarily used in such preparations, for example bodying agents, stabilizers, fillers, preservatives, perfumes, antifoams, dyes, pigments which have a coloring action, thickeners, surface-active substances, emulsifiers, emollients, moisturizers and/or humectants, anti-inflammatory substances, additional active ingredients such as vitamins or proteins, sunscreens, insect repellants,
10 bactericides, virucides, water, salts, antimicrobial, proteolytic or keratolytic substances, medicaments or other customary constituents of a cosmetic or dermatological formulation such as alcohols, polyols, polymers, foam stabilizers, organic solvents or also electrolytes.

15 The latter can be chosen, for example, from the group of salts containing the following anions: chlorides, also inorganic oxo element anions, of these, in particular sulfates, carbonates, phosphates, borates and aluminates. Electrolytes based on organic anions are also advantageous, e.g. lactates, acetates, benzoates, propionates, tartrates, citrates, amino acids, ethylenediaminetetraacetic acid and salts thereof and
20 others. Preferred cations of the salts are ammonium, alkylammonium, alkali metal, alkaline earth metal, magnesium, iron or zinc ions. It goes without saying that only physiologically acceptable electrolytes should be used in cosmetics. Particular preference is given to potassium chloride, sodium chloride, magnesium sulfate, zinc sulfate and mixtures thereof.

25 Corresponding requirements apply mutatis mutandis to the formulation of medicinal preparations.

The emulsions according to the invention can be used as a base for cosmetic or
30 dermatological formulations. The latter can have the customary composition and be used, for example, for the treatment and care of the skin and/or the hair, as lip care product, as deodorant product and as make-up or make-up remover product in decorative cosmetics or as a sunscreen preparation. For use, the cosmetic and dermatological preparations according to the invention are applied to the skin and/or
35 the hair in a sufficient amount in a manner customary for cosmetics or dermatological compositions.

For the purposes of the present invention, cosmetic or topical dermatological compositions can accordingly, depending on their composition, be used, for example, as a skin protection cream, cleansing milk, sunscreen lotion, nutrient cream, day or
5 night cream, etc. In some circumstances it is possible and advantageous to use the compositions according to the invention as a base for pharmaceutical formulations.

The cosmetic or dermatological compositions according to the invention can, for example, be in the form of preparations which can be sprayed from aerosol containers, squeezable bottles or by means of a pump device, or in the form of a liquid
10 composition which can be applied by means of roll-on devices, but also in the form of an emulsion which can be applied from normal bottles and containers.

Suitable propellants for cosmetic or dermatological preparations which can be sprayed from aerosol containers for the purposes of the present invention are the customary
15 known readily volatile, liquefied propellants, for example hydrocarbons (propane, butane, isobutane), which can be used alone or in a mixture with one another. Compressed air is also used advantageously.

20 The person skilled in the art is of course aware that there are propellants which are non-toxic per se which would be suitable in principle for realizing the present invention in the form of aerosol preparations, but which must nevertheless be avoided because of their unacceptable impact on the environment or other accompanying circumstances, in particular fluorocarbons and chlorofluorocarbons (CFCs).

25 Those cosmetic and dermatological preparations which are in the form of a sunscreen are also favorable. As well as the active ingredient combinations according to the invention, these preferably additionally comprise at least one UV-A filter substance and/or at least one UV-B filter substance and/or at least one inorganic pigment.

30 For the purposes of the present invention, however, it is also advantageous to provide those cosmetic and dermatological preparations whose main purpose is not protection against sunlight, but which nevertheless have a content of UV protectants. Thus, for example, UV-A or UV-B filter substances are usually incorporated into day creams.

35

UV protectants, like antioxidants and, if desired, preservatives, also effectively protect the preparations themselves against decay.

Furthermore, preparations according to the invention can advantageously comprise substances which absorb UV radiation in the UV-B range, the total amount of filter substances being, for example, from 0.1% by weight to 30% by weight, preferably from 0.5 to 10% by weight, in particular from 1.0 to 6.0% by weight, based on the total weight of the preparations, in order to provide cosmetic preparations which protect the hair and/or the skin from the whole region of ultraviolet radiation. They can also be used as sunscreens for the hair or the skin.

If the emulsions according to the invention comprise UV-B filter substances, the latter may be oil-soluble or water-soluble. Examples of oil-soluble UV-B filters which are advantageous according to the invention are:

- 15 - 3-benzylidenecamphor derivatives, preferably 3-(4-methylbenzylidene)camphor, 3-benzylidenecamphor;
- 4-aminobenzoic acid derivatives, preferably 2-ethylhexyl 4-(dimethylamino)benzoate, amyl 4-(dimethylamino)benzoate;
- esters of cinnamic acid, preferably 2-ethylhexyl 4-methoxycinnamate, isopentyl 4-methoxycinnamate;
- 20 - esters of salicylic acid, preferably 2-ethylhexyl salicylate, 4-isopropylbenzyl salicylate, homomenthyl salicylate;
- derivatives of benzophenone, preferably 2-hydroxy-4-methoxybenzophenone, 2-hydroxy-4-methoxy-4'-methylbenzophenone, 2,2'-dihydroxy-4-methoxybenzophenone;
- 25 - esters of benzalmalonic acid, preferably di(2-ethylhexyl) 4-methoxybenzalmalonate;
- derivatives of 1,3,5-triazine, preferably 2,4,6-trianilino(p-carbo-2'-ethyl-1'-hexyloxy)-1,3,5-triazine.

30

The list of said UV-B filters, which may be used in combination with the active ingredient combinations according to the invention, is of course not intended to be limiting.

It can also be advantageous to formulate lipodispersions according to the invention with UV-A filters which have hitherto been customarily present in cosmetic preparations. These substances are preferably derivatives of dibenzoylmethane, in

particular 1-(4'-tert-butylphenyl)-3-(4'-methoxyphenyl)propane-1,3-dione and 1-phenyl-3-(4'-isopropylphenyl)propane-1,3-dione.

Cosmetic and dermatological preparations according to the invention can also
5 comprise inorganic pigments which are customarily used in cosmetics for protecting
the skin against UV rays. These are oxides of titanium, zinc, iron, zirconium, silicon,
manganese, aluminum, cerium and mixtures thereof, and modifications in which the
oxides are the active agents. Particular preference is given to pigments based on
titanium dioxide.

10

Further constituents which can be used are:

- fats, waxes and other natural and synthetic fatty substances, preferably esters
of fatty acids with alcohols of low carbon number, e.g. with isopropanol,
propylene glycol or glycerol, or esters of fatty alcohols with alcanoic acids of
15 low carbon number or with fatty acids;
- alcohols, diols or polyols of low carbon number, and ethers thereof, preferably
ethanol, isopropanol, propylene glycol, glycerol, ethylene glycol, ethylene glycol
monoethyl or monobutyl ethers, propylene glycol monomethyl, monoethyl or
monobutyl ethers, diethylene glycol monomethyl or monoethyl ethers and
20 analogous products.

20

The examples below serve to illustrate the present invention without limiting it. The
numerical values in the examples refer to percentages by weight, based on the total
weight of the respective preparations.

25

28

Formulation examples**Example 1:**

		% by wt
	PEG-30 glyceryl stearate	2.50
5	Glycerol monostearate	1.00
	Cetyl alcohol	1.00
	Vaseline	2.50
	Polyisobutene	8.00
	Cyclomethicone	5.00
10	Aristoflex AVC	0.20
	Glycerol	5.00
	Tocopherol acetate	1.00
	Perfume, preservatives, NaOH, dyes, antioxidants etc.	q.s.
15	Water	ad 100.00

Example 2:

		% by wt
20	Glyceryl stearate citrate	2.50
	Cetyl alcohol	1.00
	Caprylic/capric triglycerides	5.00
	Cyclomethicone	5.00
	Octyldodecanol	5.00
25	Aristoflex AVC	0.30
	Glycerol	3.00
	Perfume, preservatives, NaOH, dyes, antioxidants etc.	q.s.
	Water	ad 100.00

30

29

Example 3:

	% by wt.
Stearic acid	1.50
Sorbitan monostearate	0.50
5 Myristyl alcohol	1.00
Glycerol monostearate	0.50
Paraffin oil, subliquidum	10.00
Dimethicone	1.00
Octyldodecanol	2.00
10 Hydrogenated coconut fatty acid glycerides	0.50
Aristoflex AVC	0.30
Serine	0.50
Glycerol	5.00
Tocopherol acetate	0.50
15 Perfume, preservatives, NaOH, dyes, antioxidants etc.	q.s.
Water	ad 100.00

Example 4:

	% by wt.
20 Sorbitan monostearate	2.00
Laurylmethicone copolyol	0.35
Cetylmethicone copolyol	0.15
Paraffin oil, subliquidum	10.00
25 Octyldodecanol	4.00
Hydrogenated coconut fatty acid glycerides	1.00
Cyclomethicone	1.00
Dimethicone	1.00
Aristoflex AVC	0.30
30 Glycerol	5.00
Tocopherol acetate	1.00
Perfume, preservatives, NaOH dyes, antioxidants etc.	q.s.
Water	ad 100.00

30

Example 5 (Emulsion make-up):

	% by wt.
PEG-30 stearate	2.00
Glycerol monostearate	1.00
5 Paraffin oil, subliquidum	7.00
Octyldodecanol	7.00
Isopropyl lanolate	4.00
Octyl methoxycinnamate	2.00
Butylmethoxydibenzoylmethane	1.00
10 Aristoflex AVC	0.20
Glycerol	5.00
1,3-Butylene glycol	2.00
Tocopherol acetate	1.00
Starch sodium octenyl succinate	2.50
15 Magnesium silicate	1.00
Mica	1.00
Iron oxides	1.00
Titanium dioxide	2.50
Talc	5.00
20 Perfume, preservatives, NaOH, dyes, antioxidants etc.	q.s.
Water	ad 100.00

31

Example 6 (Liquid eye liner):

	% by wt.
Stearic acid	1.20
Isopropyl lanolate	1.20
5 Dimethicone	0.40
Hydrogenated palm fatty acid glycerides	1.70
Color pigments	20.00
Aristoflex AVC	0.25
Magnesium aluminum silicate	0.30
10 1,3-Butylene glycol	4.00
Triethanolamine	0.40
Ethanol	10.00
Perfume, preservatives, antioxidants, etc.	q.s.
Water	ad 100.00
15	

Example 7 (Skin tone perfector):

	% by wt.
Color pigments	10.00
20 Cyclomethicone	25.00
Dimethicone	10.00
1,3-Butylene glycol	4.50
Glycerol	3.50
Polysorbate 40	3.50
25 Decyl oleate	2.00
Na hyaluronate	0.10
Aristoflex AVC	0.30
Perfume, preservatives, NaOH, dyes, antioxidants, etc.	q.s.
30 Water	ad 100.00

Claims:

1. A cosmetic or dermatological emulsion of the oil-in-water type, comprising
 - (i) up to 90% by weight of a water phase,
 - 5 (ii) up to 40% by weight of a lipid phase, based on the total weight of the preparation,
 - (iii) up to 10% by weight of one or more emulsifiers,
 - (iv) also comprising up to 5% by weight of one or more ammonium acryloyldimethyl-taurates/vinylpyrrolidone copolymers.
- 10 2. The emulsion as claimed in claim 1, wherein its lipid content is chosen from the range from 0.5% by weight to 20% by weight, preferably from 5 to 10% by weight.
3. The emulsion as claimed in claim 1, wherein its lipid content is up to 7.5% by
- 15 weight.
4. The emulsion as claimed in claim 1, which comprises one or more dyes and/or coloring pigments.
- 20 5. The emulsion as claimed in claim 4, wherein the total amount of the dyes and coloring pigments is chosen from the range from, for example, 0.1% by weight to 30% by weight, preferably from 0.5 to 15% by weight, in particular from 1.0 to 10% by weight, in each case based on the total weight of the preparations.

33

Abstract:

Cosmetic or dermatological gel creams of the oil-in-water type, comprising

- (i) up to 90% by weight of a water phase,
- 5 (ii) up to 20% by weight of a lipid phase, based on the total weight of the preparations,
- (iii) up to 5% by weight of one or more emulsifiers,
- (iv) also comprising up to 5% by weight of one or more ammonium acryloyldimethyl-taurates/vinylpyrrolidone copolymers.

10

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.